

## **Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players**

Ramírez-Campillo, Rodrigo; González-Jurado, José Antonio; Martínez, Cristian; Nakamura, Fábio Yuzo; Peñailillo, Luis; Meylan, Cesar M.P.; Caniuqueo, Alexis; Cañas-Jamet, Rodrigo; Moran, Jason; Alonso-Martínez, Alicia M.; Izquierdo, Mikel

*Published in:*

Journal of Science and Medicine in Sport

*Publication date:*

2016

*The re-use license for this item is:*

CC BY-NC-ND

*This document version is the:*

Peer reviewed version

*The final published version is available direct from the publisher website at:*  
[10.1016/j.jsams.2015.10.005](https://doi.org/10.1016/j.jsams.2015.10.005)

**Find this output at Hartpury Pure**

*Citation for published version (APA):*

Ramírez-Campillo, R., González-Jurado, J. A., Martínez, C., Nakamura, F. Y., Peñailillo, L., Meylan, C. M. P., Caniuqueo, A., Cañas-Jamet, R., Moran, J., Alonso-Martínez, A. M., & Izquierdo, M. (2016). Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players. *Journal of Science and Medicine in Sport*, 19(8), 682-687. <https://doi.org/10.1016/j.jsams.2015.10.005>

## Accepted Manuscript

Title: Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players

Author: Rodrigo Ramírez-Campillo José Antonio  
González-Jurado Cristian Martínez Fábio Yuzo Nakamura  
Luis Peñailillo Cesar M.P. Meylan Alexis Caniuqueo Rodrigo  
Cañas-Jamet Jason Moran Alicia M. Alonso-Martínez Mikel  
Izquierdo



PII: S1440-2440(15)00205-4  
DOI: <http://dx.doi.org/doi:10.1016/j.jsams.2015.10.005>  
Reference: JSAMS 1237

To appear in: *Journal of Science and Medicine in Sport*

Received date: 24-8-2015  
Revised date: 25-9-2015  
Accepted date: 29-10-2015

Please cite this article as: Ramírez-Campillo R, González-Jurado JA, Martínez C, Nakamura FY, Peñailillo L, Meylan CMP, Caniuqueo A, Cañas-Jamet R, Moran J, Alonso-Martínez AM, Izquierdo M, Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players, *Journal of Science and Medicine in Sport* (2015), <http://dx.doi.org/10.1016/j.jsams.2015.10.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Effects of plyometric training and creatine supplementation on maximal-intensity exercise and**  
2 **endurance in female soccer players**

3  
4 Rodrigo Ramírez-Campillo <sup>a,b</sup>, José Antonio González-Jurado <sup>c</sup>, Cristian Martínez <sup>d</sup>, Fábio Yuzo  
5 Nakamura <sup>e</sup>, Luis Peñailillo <sup>f</sup>, Cesar M.P. Meylan <sup>g,h</sup>, Alexis Caniuqueo <sup>i</sup>, Rodrigo Cañas-Jamet <sup>j,k</sup>, Jason  
6 Moran <sup>l</sup>, Alicia M. Alonso-Martínez <sup>m</sup>, Mikel Izquierdo <sup>m</sup>

7  
8 <sup>a</sup> Department of Physical Activity Sciences, Universidad de Los Lagos, Osorno, Chile  
9 (r.ramirez@ulagos.cl)

10 <sup>b</sup> Laboratory of Exercise Sciences, MEDS Clinic, Santiago, Chile

11 <sup>c</sup> Faculty of Sport, University Pablo de Olavide, Sevilla, Spain (jagonjur@upo.es)

12 <sup>d</sup> Department of Physical Education, Sport and Recreation, Universidad de La Frontera, Temuco, Chile  
13 (cristian.martinez.s@ufrontera.cl)

14 <sup>e</sup> Departamento de Educação Física, Universidade Estadual de Londrina, Londrina, Brazil  
15 (fabioy\_nakamura@yahoo.com.br)

16 <sup>f</sup> Exercise Science Laboratory, School of Kinesiology, Faculty of Medicine, Universidad Finis Terrae,  
17 Santiago, Chile (lpenailillo@uft.cl)

18 <sup>g</sup> Canadian Sport Institute Pacific, Vancouver, Canada (cmeylan@csipacific.ca)

19 <sup>h</sup> Canadian Soccer Association, Ottawa, Canada

<sup>i</sup> Laboratory of Physiology and Biomechanics, Universidad Autónoma de Chile, Temuco, Chile  
(alexis.caniuqueo@uaautonoma.cl)

20 <sup>j</sup> Laboratorio de Fisiología, Departamento de Ciencias Biológicas, Facultad de Ciencias Biológicas,  
21 Universidad Andrés Bello, Viña del Mar, Chile (rcanaskine@gmail.com)

22 <sup>k</sup> Escuela de Kinesiología, Facultad de Medicina, Universidad Diego Portales, Santiago, Chile.

23 <sup>l</sup> Centre for Sports and Exercise Science, School of Biological Sciences, University of Essex, Colchester,  
24 United Kingdom (jmorana@essex.ac.uk)

25 <sup>m</sup> Department of Health Sciences, Public University of Navarre, Navarra, Spain  
26 (mikel.izquierdo@gmail.com)

27

28

29 **Corresponding author**

30 Mikel Izquierdo, PhD

31 Department of Health Sciences

32 Public University of Navarre (Navarra) SPAIN

33 Campus of Tudela

34 Av. de Tarazona s/n. 31500 Tudela (Navarra) SPAIN

35 Tel + 34 948 417876

36 mikel.izquierdo@gmail.com

37

38 **Word count:** 2690

41 **Abstract**

42 *Objectives:* to investigate the effects of a six-week plyometric training and creatine supplementation  
43 intervention on maximal-intensity and endurance performance in female soccer players during in-season  
44 training.

45 *Design:* Randomized, double-blind, placebo-controlled trial

46 *Methods:* Young (age  $22.9 \pm 2.5$  y) female players with similar training load and competitive background  
47 were assigned to a plyometric training group receiving placebo (PLACEBO, n = 10), a plyometric training  
48 group receiving creatine supplementation (CREATINE, n = 10) or a control group receiving placebo  
49 without following a plyometric program (CONTROL, n = 10). Athletes were evaluated for jumping,  
50 maximal and repeated sprinting, endurance and change-of-direction speed performance before and after  
51 six weeks of training.

52 *Results:* After intervention the CONTROL group did not change, whereas both plyometric training groups  
53 improved jumps (ES = 0.25-0.49), sprint (ES = 0.35-0.41), repeated sprinting (ES = 0.48-0.55), endurance  
54 (ES = 0.32-0.34) and change-of-direction speed performance (ES = 0.46-0.55). However, the CREATINE  
55 group improved more in the jumps and repeated sprinting performance tests than the CONTROL and the  
56 PLACEBO groups.

57 *Conclusions:* Adaptations to plyometric training may be enhanced with creatine supplementation.

58

59 **Key words:** muscle strength; sports; women; strength training; ergogenic aids.

60

61

62

63

64

65

66

## 67 1. Introduction

68 Soccer players must perform numerous single maximal-intensity exercises, including jumping,  
69 kicking, accelerating and decelerating<sup>1</sup>, actions that might precede most of the goals scored in competitive  
70 leagues<sup>2</sup>, and correlate with competition success<sup>3</sup>. Repeating these maximal-intensity actions across a 90-  
71 min game is also important<sup>4</sup> and might be associated with endurance<sup>5</sup>, but also with intramuscular creatine  
72 phosphate<sup>6</sup>, a critical energy source for maximal-intensity actions. Therefore, investigating the methods by  
73 which single and repeated maximal-intensity actions (alongside endurance) can be enhanced in female  
74 soccer players is important for this population. Plyometric training in female players may improve their  
75 maximal-intensity exercise and endurance<sup>7</sup>. However, further investigation in this population is required<sup>8</sup>,  
76 especially in regard to factors that might be mediating the effects of plyometric training on maximal-  
77 intensity exercise and endurance performances adaptations, such as dietary supplements<sup>9</sup>.

78 Previous research involving male<sup>10</sup> as well as female soccer players<sup>11</sup> has demonstrated that acute  
79 creatine intake (i.e., one week) can enhance maximal-intensity exercise (e.g., jump, sprint, agility).  
80 Despite these meaningful results, recently it has been shown that acute creatine supplementation had no  
81 positive effects on fatigue and repeated sprint ability in a match simulation protocol<sup>12</sup>, suggesting that  
82 longer-term use might be more beneficial to performance<sup>13</sup>. Among the few longitudinal studies conducted  
83 with regard to soccer, during a seven-week functional overreaching pre-season, creatine supplementation  
84 prevented deterioration of male soccer players' maximal-intensity performance<sup>14</sup>. In female players,  
85 creatine showed a positive effect on strength during a 13-week off-season<sup>15</sup>. However, to our knowledge,  
86 it remains unknown whether creatine supplementation and plyometric training can elicit similar  
87 improvements in female players when compared to plyometric training alone.

88 Therefore, the objective of this study was to investigate the effects of a six-week plyometric  
89 training and creatine supplementation intervention on maximal-intensity and endurance performance in  
90 female soccer players.

91

92

## 93 2. Methods

94 After written informed consent, 33 amateur female players (three goalkeepers, nine defenders, ten  
95 midfielders and eleven forwards) participated in this study. Participants had no regular strength or  
96 plyometric training during the three months prior to the intervention and had never before taken creatine  
97 supplements. The sample size was determined according to changes in vertical jump performance in a  
98 group of soccer players submitted to a control ( $\Delta = 0.5$  cm;  $SD = 1.1$ ) or to a short-term plyometric  
99 training ( $\Delta = 2.6$  cm;  $SD = 1.6$ )<sup>16</sup> comparable with that performed in this study. Eight participants per  
100 group would yield a power of 95% and  $\alpha = 0.01$ , with a detectable ES of 0.2.

101 Exclusion criteria included (a) potential medical problems that compromised participation or  
102 performance in the study, (b) any lower-extremity surgery in the past two years, (c) previous use creatine  
103 consumption. Based on these criteria, three participants were excluded (one defender and two midfielders  
104 were identified with recent history of ankle or knee injury). The participants included in the study  
105 (between 19 and 28 y of age) were randomly assigned to either a plyometric training group receiving  
106 placebo (PLACEBO,  $n = 10$ ), a plyometric training group receiving creatine supplementation  
107 (CREATINE,  $n = 10$ ) or a control group receiving placebo without following a plyometric program  
108 (CONTROL,  $n = 10$ ). No vegetarians were registered in the study. At baseline, no differences were  
109 observed in any descriptive (or dependent) variable between groups (Table 1). The study was conducted in  
110 accordance with the Declaration of Helsinki and was approved by the ethics committee of the responsible  
111 department.

112  
113 **\*\*\*Table 1 near here\*\*\***  
114

115 Participants were accustomed to the testing procedures, as athletes incorporate them as a regular  
116 aspect of their training schedule. Measurements were taken one week before and after intervention,  
117 completed in three non-consecutive days and were always administered in the same order, at the same

118 time of the day and by the same investigators, who were blinded to each participant's group assignment.  
119 Ten minutes of standard warm-up was performed before testing.

120 On day one, height, body mass, squat jump, countermovement jump, 20-m sprint test and running  
121 anaerobic sprint test (RAST) measurements were completed. On day two, 20-cm and 40-cm drop jump  
122 reactive strength indexes, peak jump power and peak jump power load testing were completed. On day  
123 three, unilateral 20-cm drop jump reactive strength indexes (right and left leg), change-of-direction speed  
124 test (i.e., *Illinois* test) and 20-m multi stage shuttle run tests were completed. Three maximal trials were  
125 allowed for all performance tests, excepting the single shuttle run test, peak jump power test and RAST  
126 measurements. At least two minutes of rest were permitted between each maximal trial to reduce the  
127 effects of fatigue.

128 Anthropometric measurements employed a stadiometer (Bodometer 206, SECA, Hamburg,  
129 Germany) and an electrical scale (BF 100\_Body Complete, BEURER, Ulm, Germany). Test protocols for  
130 the jumps, 20-m sprints, change-of-direction speed<sup>17</sup> and shuttle run tests<sup>18</sup> were performed as previously  
131 described. Briefly, for the jumps, players executed maximal effort jumps on a mobile contact mat  
132 (Ergojump; Globus, Codogne, Italy) with arms akimbo. Take-off and landing were standardized to full  
133 knee and ankle extension on the same spot. The participants were instructed to maximize jump height. In  
134 addition, for the 20 and 40 cm drop jump reactive strength index, players were instructed to minimize  
135 ground contact time after dropping down from a 20 and 40 cm drop box, respectively. For the 20-m  
136 sprints, participants had a standing start with the toe of the preferred foot forward and just behind the  
137 starting line. For the change of direction speed test, the timing system and procedures were same as for the  
138 20-m sprint test, except that players started supine and completed a circuit with several changes of  
139 directions. For the shuttle run endurance test, players ran back and forth between two lines, spaced 20-m  
140 apart, in time with the "beep" sounds from an electronic audio recording. Each successful run of the 20-m  
141 distance was a completion of a shuttle. The beep sounded at a progressively increasing pace with every  
142 minute of the test, and the player had to increase speed accordingly until volitional fatigue. For unilateral

143 jumps, instead of using both legs during jumping and landing, participants used their left and right legs  
144 alternatively.

145 Peak jump power measurements employed the same equipment and movement patterns as  
146 countermovement jump measurements; however, instead of adopting arms akimbo, participants put weight  
147 bars on their shoulders. To estimate power (W), a previously established testing protocol<sup>19</sup> and equation<sup>20</sup>  
148  $[W = 65.1 \times \text{jump height (cm)} \times 25.8 \times \text{body mass (kg)} - 1413.1]$  was used. Briefly, unloaded peak jump  
149 power was determined with a broomstick, while, in the following attempts, loads were increased by 5 kg  
150 and tests were stopped when reductions in power output were greater than 50 W compared to previous  
151 jump load measurements. Though the number of attempts were not predetermined, >90% of athletes  
152 completed between four and six attempts, reducing the probability that fatigue affected test outcomes.

153 Participants performed six 35-m maximal sprints with 10 s of rest for the RAST, as previously  
154 described and validated elsewhere<sup>21</sup>. The start for each sprint (10-s interval) occurred with a sound from  
155 the measurement equipment. Sprint times were measured using single beam infrared photoelectric cells  
156 (Globus Italy, Codogne, Italy) leveled ~0.7 m above the floor (i.e., hip level). The starting position was  
157 standardized to a split position with the toe of each preferred foot forward and behind the starting line.  
158 Mean RAST times were used for analyses.

159 All groups participated in the same soccer training program, such that similar training loads were  
160 measured by session rating of perceived exertion (RPE), as previously described<sup>7</sup> (Table 1). Briefly, each  
161 player's session RPE was collected ~30 min after each soccer training session and match to ensure that the  
162 perceived effort reflected the entire session rather than the most recent exercise intensity. Total training  
163 load was calculated as RPE  $\times$  training session duration (i.e., minutes).

164 Experiments were completed during competition (i.e., mid portion of the in-season), which was  
165 similar between groups (Table 1). Participants in the plyometric training groups performed plyometric  
166 drills immediately after warm-up and as a substitute for some soccer drills (i.e., technical-tactical) within  
167 the usual 120-minute practice twice per week for six weeks. Plyometric intervention was determined  
168 based on previous research regarding soccer players<sup>16</sup>. A detailed description of the training program can



169 be found in a previous study<sup>18</sup>. Briefly, plyometric training included unilateral and bilateral horizontal and  
170 vertical jumps with both cyclic and acyclic arm swings. Participants were motivated to achieve maximal  
171 effort in every jump, instructed to aim toward maximal vertical heights and horizontal distances for  
172 acyclic jumps and minimum ground contact times for cyclic jumps, in order to maximize reactive strength.

173 Before the training period, participants were accustomed to all exercises completed in the  
174 plyometric program, and all training sessions were supervised with a coach to player ratio of 1:3, with  
175 particular attention paid to technique. Plyometric training sessions were separated by a minimum of 48  
176 hours (including games). Each plyometric training group completed the same number of total jumps, with  
177 the same progressive overload, used the same surface and time of day for training and the same rest  
178 intervals between jumps (i.e., 15 s for acyclic jumps) and sets (i.e., 60 s).

179 The CREATINE group participants received 20 g/d of creatine monohydrate (Gnc Pro  
180 Performance, USA), divided into four equal doses, over the course of one week, followed by single daily  
181 doses of 5 g for the next five weeks<sup>14</sup>. Participants in the PLACEBO and CONTROL groups were given  
182 the same dosages of glucose. During the loading phase, supplements were presented in four packages, and  
183 participants were instructed to ingest the packet contents at breakfast, lunch, dinner and before bedtime.  
184 During the maintenance phase, each participant consumed the supplement as a single dose during her  
185 lunch. To mask the taste and texture of the supplements provided to them, participants were asked to  
186 dissolve the supplements in juice that contained a small amount of carbohydrates to reduce creatine  
187 muscle uptake. Compliance to supplementation was monitored weekly via personal communication. Only  
188 one athlete in the CREATINE group reported mild gastrointestinal distress, but this participant completed  
189 the study. The supplement packages were coded, so that neither the investigators nor the participants were  
190 aware of the contents until completion of the analyses. The supplements were distributed by a staff  
191 member who was not an investigator in this study. Participants' feedback on group assignments post study  
192 demonstrated the effectiveness of the double blinded protocol (in which 30% of participants guessed their  
193 group assignments).

194 One week immediately before and after intervention, each participant's energy, macronutrient and  
195 creatine intakes were determined through a 24-hour food recall questionnaire conducted in three different  
196 days of the week, as previously described<sup>22</sup>.

197 Statistical analyses employed the STATISTICA statistical package (Version 8.0; StatSoft, Inc,  
198 Tulsa). All values are reported as the means  $\pm$  standard deviations. Relative changes (%) in performance  
199 and Cohen's d-effect sizes (ES) are expressed with 90% confidence limits. Normality and  
200 homoscedasticity assumptions made for all data before and after intervention were checked using the  
201 Shapiro-Wilk and Levene tests, respectively. To determine the effects of the intervention on performance  
202 adaptations, groups were compared using mixed-design factorial ANOVA. When a significant F value  
203 occurred for interaction between groups or for main effects of group or time, Tukey post hoc procedures  
204 were performed. In addition, a between-groups one-way analysis of variance compared changes between  
205 groups (i.e., the differences between scores before and after the intervention). The  $\alpha$  level was set at  $p <$   
206 0.05 for statistical significance. In addition to this null hypothesis testing, data were also assessed for  
207 practical meaningfulness using a magnitude-based inference approach. Threshold values for assessing  
208 magnitudes of ES were 0.20, 0.60, 1.2, and 2.0 for small, moderate, large, and very large, respectively<sup>23</sup>.  
209 Magnitudes of differences in training effects between groups were evaluated non-clinically<sup>23</sup>: if the  
210 confidence interval overlapped thresholds for substantial positive and negative values, the effect was  
211 deemed unclear (i.e., trivial). The effect was otherwise clear and reported as the magnitude of the  
212 observed value with a qualitative probability, as above (i.e., small, moderate, large, and very large).

213

### 214 3. Results

215 The reliability of assessments was determined using the typical error of measurement expressed as  
216 a percentage of the mean (i.e., coefficient of variation) and ranged from 0.8 to 5.8%.

217 The energy, carbohydrate, lipids, protein and creatine intakes did not differ before, during and  
218 after the intervention for the CONTROL ( $2678 \pm 427$  kcal·day<sup>-1</sup>;  $377 \pm 89.8$  g·day<sup>-1</sup>;  $91.1 \pm 23.8$ ;  $88.1 \pm$   
219  $25.4$ ;  $1.2 \pm 0.5$  g·day<sup>-1</sup>, respectively), PLACEBO ( $2819 \pm 242$  kcal·day<sup>-1</sup>;  $420 \pm 61.2$  g·day<sup>-1</sup>;  $86.1 \pm 10.9$

220  $\text{g}\cdot\text{day}^{-1}$ ;  $91.4 \pm 15.3 \text{ g}\cdot\text{day}^{-1}$ ;  $1.2 \pm 0.4 \text{ g}\cdot\text{day}^{-1}$ , respectively) or CREATINE group ( $2635 \pm 325 \text{ kcal}\cdot\text{day}^{-1}$ ;  
221  $383 \pm 66.4 \text{ g}\cdot\text{day}^{-1}$ ;  $84.2 \pm 15.9 \text{ g}\cdot\text{day}^{-1}$ ;  $86.3 \pm 10.9 \text{ g}\cdot\text{day}^{-1}$ ;  $1.3 \pm 0.4 \text{ g}\cdot\text{day}^{-1}$ , respectively). Similarly,  
222 body mass and body mass index were not different before, during and after the intervention for the  
223 CONTROL ( $60.1 \pm 7.5 \text{ kg}$ ;  $23.3 \pm 2.2 \text{ kg}\cdot\text{m}^{-2}$ , respectively) or PLACEBO ( $56.8 \pm 5.4 \text{ kg}$ ;  $21.2 \pm 1.4 \text{ kg}\cdot\text{m}^{-2}$ ,  
224  $^2$ , respectively) groups. However, regarding the basal value of body mass ( $60.4 \pm 8.0 \text{ kg}$ ) and body mass  
225 index ( $23.2 \pm 3.1 \text{ kg}\cdot\text{m}^{-2}$ ) of the CREATINE group, an increase ( $p < 0.05$ ; 1.4%) was observed during the  
226 experimental period.

227 Both plyometric training groups (CREATINE and PLACEBO) increased ( $p < 0.05$ ) jump and  
228 power performance (ES = 0.23-0.49), however, only the CREATINE group showed a greater increase  
229 compared with the CONTROL group (Table 2). In addition, the CREATINE group had small greater  
230 meaningful training effects on peak jump power load, squat jump and 40-cm drop jump reactive strength  
231 index compared to PLACEBO group (Table 2).

232  
233 **\*\*\*Table 2 around here\*\*\***  
234  
235  
236

237 Regarding RAST, change of direction speed, 20-m sprint and shuttle run endurance, both  
238 plyometric training groups increased ( $p < 0.05$ ) performance in these test (ES = 0.32-0.55), however, the  
239 CREATINE group had small greater meaningful training effects on the RAST compared to PLACEBO  
240 group and CONTROL group (Table 3).

241  
242  
243  
244 **\*\*\*Table 3 around here\*\*\***  
245

246

247 **4. Discussion**

248 Our results suggest that replacement of some soccer drills with specific plyometric training, with  
249 no additional training time during (in-season) competition, is an effective training strategy for increasing  
250 maximal-intensity and endurance performance in female soccer players. Furthermore, our results  
251 demonstrated that creatine supplementation during plyometric training may boost further adaptations  
252 related to maximal-intensity exercise and repeated sprint ability.

253 Considering that neither group changed dietary intake during the experimental period, the increase  
254 in body mass (1.4%) and body mass index in the CREATINE group could be attributed to the acute effect  
255 of creatine supplementation, as similar increases in body mass have been shown after seven days of  
256 supplementation in female soccer players (0.8%)<sup>10</sup>. Alternatively, creatine supplementation has been  
257 shown to increase the cross-sectional areas of both type I and type II muscle fibers as well as several  
258 myogenic regulatory factors during long-term training<sup>24</sup>, although this effect may be more elusive in  
259 female players after short-term training interventions<sup>15</sup>.

260 Our results indicated that both plyometric training groups improved sprint performances, change-  
261 of-direction speeds and endurance at the end of the interventions; on the other hand, no changes were  
262 observed in the CONTROL group. These results are similar to those previously reported in female soccer  
263 players<sup>7</sup>. The improvements observed after plyometric training in unidirectional (i.e., sprint)<sup>25</sup> or maximal-  
264 intensity change-of-direction maneuvers<sup>26</sup> may have been mediated by rapid (i.e., < 6-weeks)  
265 neuromuscular adaptations of targeted muscle groups<sup>27</sup>, which occur during even the most competitive  
266 periods of the athletes' calendar (i.e., in-season)<sup>7</sup>. The observed improvements in endurance might have  
267 occurred by means of cardiovascular (i.e., VO<sub>2</sub>max)<sup>28</sup>, neuromuscular-mediated changes in the athletes'  
268 running economies<sup>29</sup> or neuromuscular power improvements that affect the athletes' change-of-direction  
269 endurance results<sup>30</sup>.

270 Although both plyometric training groups increased their sprint performances, change-of-direction  
271 speeds and endurance, only the CREATINE group showed a meaningful increase in peak jump power

272 load (Table 2). More so, the CREATINE group showed a greater increase in 40-cm drop jump reactive  
273 strength index, peak jump power load and squat jump performance compared with the CONTROL and  
274 PLACEBO groups (Table 2). The greater enhancement in muscular capabilities observed with regard to  
275 loads (i.e., peak jump power loads) and in concentric-only maximal-intensity performances (i.e., squat  
276 jumps) in the CREATINE group may be indicative of greater force-related adaptations<sup>19</sup>. Alternatively,  
277 compared to the PLACEBO group, the greater improvements in jump performance observed in the  
278 CREATINE group might have been caused by smaller decays in drills-intensity during training sessions<sup>10</sup>  
279 and reduced time required for recovery after training<sup>13</sup>. This is partially supported by the greater jump  
280 heights, jump lengths and reactive strength indexes values (used as biofeedback in some training sessions)  
281 observed (although not reported in this manuscript) in the CREATINE group during plyometric training  
282 sessions.

283 RAST mean sprint times improved for both PLACEBO and CREATINE groups after plyometric  
284 training (Table 3). However, compared to the PLACEBO group, the CREATINE group showed greater  
285 improvement in RAST mean sprint times (Table 3). To the author's knowledge, the results reported herein  
286 are novel, in that our experiments involved female soccer players, making comparisons with previous  
287 studies difficult. However, creatine supplementation have shown to increase repeated sprint abilities of  
288 male soccer players<sup>10</sup>, possible by means of increasing phosphocreatine re-synthesis rates during  
289 recoveries between RAST sprints<sup>6</sup>; this adaptation might help explain the greater RAST improvements  
290 observed in the CREATINE group compared to the PLACEBO group. As sprints<sup>1</sup> and their repetition<sup>4</sup>  
291 during games are key aspects of soccer competition, and might be related with goals scored<sup>2</sup> and success<sup>3</sup>,  
292 the greater power during repeated sprints<sup>21</sup> achieved by the CREATINE group might allow female players  
293 to achieve an important competitive advantage.

294

## 295 **5. Conclusions**

296 For female soccer players, replacement of some low-intensity technical-tactical soccer drills  
297 during the in-season period with maximal-intensity exercise plyometric drills, in a short-term (i.e., 6

298 weeks) plyometric training intervention, induced higher maximal-intensity exercise and endurance  
299 performance improvements compared to soccer training alone, and the improvements induced by  
300 plyometric training were enhanced by creatine supplementation. In practical terms, creatine  
301 supplementation may be seen as an ergogenic aid while applying plyometric training in adult female  
302 soccer players, at least when the target is improving specific physical performance.

303

### 304 **Practical Implications**

305 · Replacing some low-intensity technical-tactical soccer drills with plyometric drills might induce higher  
306 maximal-intensity and endurance performance improvements in participating female soccer players,  
307 compared to soccer-training alone.

308 · Maximal-intensity and endurance performance improvements induced by plyometric drills might be  
309 enhanced by creatine supplementation, particularly in task where a shift in force production might result in  
310 more powerful movements (i.e., loads at peak power or SJ) or where increases in intramuscular creatine  
311 content are relevant (i.e., RAST).

312 · Creatine supplementation, when combined with plyometric training, show no detrimental effect on  
313 endurance performance of female soccer players during a short-term in-season competitive period.

314

### 315 **Acknowledgements**

316 The authors wish to thank all the volunteers who participated in this study. There was no financial  
317 assistance with this project.

318

### 319 **References**

- 320 1. Barnes C, Archer DT, Hogg B et al. The evolution of physical and technical performance  
321 parameters in the English Premier League. *Int J Sports Med* 2014; 35(13):1095-1100.
- 322 2. Faude O, Koch T, Meyer T. Straight sprinting is the most frequent action in goal situations in  
323 professional football. *J Sports Sci* 2012; 30(7):625-631.

- 324 3. Arnason A, Sigurdsson SB, Gudmundsson A et al. Physical fitness, injuries, and team  
325 performance in soccer. *Med Sci Sports Exerc* 2004; 36(2):278-285.
- 326 4. Carling C, Le Gall F, Dupont G. Analysis of repeated high-intensity running performance in  
327 professional soccer. *J Sports Sci* 2012; 30(4):325-336.
- 328 5. Helgerud J, Engen LC, Wisloff U et al. Aerobic endurance training improves soccer performance.  
329 *Med Sci Sports Exerc* 2001; 33(11):1925-1931.
- 330 6. Girard O, Mendez-Villanueva A, Bishop D. Repeated-sprint ability - part I: factors contributing to  
331 fatigue. *Sports Med* 2011; 41(8):673-694.
- 332 7. Ramirez-Campillo R, Vergara-Pedrerros M, Henriquez-Olguin C et al. Effects of plyometric  
333 training on maximal-intensity exercise and endurance in male and female soccer players. *J Sports*  
334 *Sci*. In press. doi: 10.1080/02640414.2015.1068439
- 335 8. Datson N, Hulton A, Andersson H et al. Applied physiology of female soccer: an update. *Sports*  
336 *Med* 2014; 44(9):1225-1240.
- 337 9. Hespel P, Maughan RJ, Greenhaff PL. Dietary supplements for football. *J Sports Sci* 2006;  
338 24(7):749-761.
- 339 10. Mujika I, Padilla S, Ibanez J et al. Creatine supplementation and sprint performance in soccer  
340 players. *Med Sci Sports Exerc* 2000; 32(2):518-525.
- 341 11. Cox G, Mujika I, Tumilty D et al. Acute creatine supplementation and performance during a field  
342 test simulating match play in elite female soccer players. *Int J Sport Nutr Exerc Metab* 2002;  
343 12(1):33-46.
- 344 12. Williams J, Abt G, Kilding AE. Effects of creatine monohydrate supplementation on simulated  
345 soccer performance. *Int J Sports Physiol Perform* 2014; 9(3):503-510.
- 346 13. Rosene J, Matthews T, Ryan C et al. Short and longer-term effects of creatine supplementation on  
347 exercise induced muscle damage. *J Sports Sci Med* 2009; 8(1):89-96.
- 348 14. Claudino JG, Mezencio B, Amaral S et al. Creatine monohydrate supplementation on lower-limb  
349 muscle power in Brazilian elite soccer players. *J Int Soc Sports Nutr* 2014; 11:32.

- 350 15. Larson-Meyer DE, Hunter GR, Trowbridge CA et al. The effect of creatine supplementation on  
351 muscle strength and body composition during off-season training in female soccer players. *J*  
352 *Strength Cond Res* 2000; 14(4):434-442.
- 353 16. Ramirez-Campillo R, Burgos C, Henriquez-Olguin C et al. Effect of unilateral, bilateral and  
354 combined plyometric training on explosive and endurance performance of young soccer players. *J*  
355 *Strength Cond Res* 2015; 29(5):1317-1328.
- 356 17. Ramirez-Campillo R, Andrade DC, Izquierdo M. Effects of plyometric training volume and  
357 training surface on explosive strength. *J Strength Cond Res* 2013; 27(10):2714-2722.
- 358 18. Ramirez-Campillo R, Meylan CM, Alvarez-Lepin C et al. The effects of interday rest on  
359 adaptation to 6-weeks of plyometric training in young soccer players. *J Strength Cond Res* 2015;  
360 29(4):972-979.
- 361 19. Jimenez-Reyes P, Samozino P, Cuadrado-Penafiel V et al. Effect of countermovement on power-  
362 force-velocity profile. *Eur J Appl Physiol* 2014; 114(11):2281-2288.
- 363 20. Canavan PK, Vescovi JD. Evaluation of power prediction equations: peak vertical jumping power  
364 in women. *Med Sci Sports Exerc* 2004; 36(9):1589-1593.
- 365 21. Zagatto AM, Beck WR, Gobatto CA. Validity of the running anaerobic sprint test for assessing  
366 anaerobic power and predicting short-distance performances. *J Strength Cond Res* 2009;  
367 23(6):1820-1827.
- 368 22. Ramirez-Campillo R, Andrade DC, Campos-Jara C et al. Regional fat changes induced by  
369 localized muscle endurance resistance training. *J Strength Cond Res* 2013; 27(8):2219-2224.
- 370 23. Hopkins WG, Marshall SW, Batterham AM et al. Progressive statistics for studies in sports  
371 medicine and exercise science. *Med Sci Sports Exerc* 2009; 41(1):3-13.
- 372 24. Willoughby DS, Rosene JM. Effects of oral creatine and resistance training on myogenic  
373 regulatory factor expression. *Med Sci Sports Exerc* 2003; 35(6):923-929.
- 374 25. Chimera NJ, Swanik KA, Swanik CB et al. Effects of plyometric training on muscle-activation  
375 strategies and performance in female athletes. *J Athl Train* 2004; 39(1):24-31.



- 376 26. Zebis MK, Bencke J, Andersen LL et al. The effects of neuromuscular training on knee joint  
377 motor control during sidcutting in female elite soccer and handball players. *Clin J Sport Med*  
378 2008; 18(4):329-337.
- 379 27. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity  
380 plyometric training. *Sports Med* 2010; 40(10):859-895.
- 381 28. Grieco CR, Cortes N, Greska EK et al. Effects of a combined resistance-plyometric training  
382 program on muscular strength, running economy, and Vo<sub>2</sub>peak in division I female soccer  
383 players. *J Strength Cond Res* 2012; 26(9):2570-2576.
- 384 29. Yamamoto LM, Lopez RM, Klau JF et al. The effects of resistance training on endurance distance  
385 running performance among highly trained runners: a systematic review. *J Strength Cond Res*  
386 2008; 22(6):2036-2044.
- 387 30. Buchheit M. The 30-15 intermittent fitness test: accuracy for individualizing interval training of  
388 young intermittent sport players. *J Strength Cond Res* 2008; 22(2):365-374.

389

390

391

392

393

394

395

396

397

398

399

400

401

402  
403  
404  
405

**Table 1.** Descriptive data of the control group (CONTROL, n = 10), plyometric training group receiving placebo (PLACEBO, n = 10) and plyometric training group receiving creatine supplementation (CREATINE, n = 10).

	CONTROL	PLACEBO	CREATINE
<b>Age (y)</b>	22.5 ± 2.1	22.9 ± 1.7	23.1 ± 3.4
<b>Body mass (kg)</b>	60.1 ± 7.5	56.8 ± 5.4	60.4 ± 8.0
<b>Height (m)</b>	1.61 ± 0.06	1.64 ± 0.09	1.62 ± 0.04
<b>Body mass index (kg.m<sup>-2</sup>)</b>	23.3 ± 2.2	21.2 ± 1.4	23.2 ± 3.1
<b>Session rating of perceived exertion<sup>a</sup></b>	468 ± 332	396 ± 234	424 ± 229
<b>Soccer experience (y)</b>	7.9 ± 3.7	7.5 ± 4.2	8.3 ± 4.7
<b>Competition games during experimental period</b>	4.0 ± 1.6	4.2 ± 1.3	4.0 ± 1.5
<b>Weekly participation in other sport or training modality (h)</b>	1.3 ± 1.1	1.2 ± 1.0	1.4 ± 1.0

Notes: <sup>a</sup>Soccer training load was determined by multiplying the minutes of soccer training by the rating of perceived exertion after each soccer training session.

406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416

417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441

**Table 2.** Training effects (with 90% confidence limits) for the jump performance variables for the control group (CONTROL, n = 10), plyometric training group receiving placebo (PLACEBO, n = 10) and plyometric training group receiving creatine supplementation (CREATINE, n = 10).

	Baseline Mean $\pm$ SD	Change (%)	Effect sizes
<b>Peak jump power (W)</b>			
CONTROL	1979 $\pm$ 211	1.7 (-2.4, 6.0)	0.13 (-0.19, 0.45)
PLACEBO	1940 $\pm$ 338	5.0 (3.6, 6.4) <sup>a</sup>	0.25 (0.18, 0.32)
CREATINE	1969 $\pm$ 250	7.0 (4.8, 9.3) <sup>b, c</sup>	0.49 (0.33, 0.64)
<b>Peak jump power load (kg)</b>			
CONTROL	14.1 $\pm$ 6.9	-8.5 (-25.8, 12.8)	-0.09 (-0.31, 0.13)
PLACEBO	14.7 $\pm$ 9.2	11.5 (-4.0, 29.6)	0.18 (-0.07, 0.42)
CREATINE	14.5 $\pm$ 8.8	20.4 (7.1, 35.4) <sup>a</sup>	0.34 (0.12, 0.55) <sup>d</sup>
<b>Squat jump (cm)</b>			
CONTROL	23.5 $\pm$ 4.1	-0.7 (-4.1, 2.9)	-0.04 (-0.22, 0.15)
PLACEBO	25.0 $\pm$ 4.5	5.1 (2.8, 7.5) <sup>a</sup>	0.27 (0.15, 0.39)
CREATINE	24.9 $\pm$ 4.4	8.3 (5.1, 11.5) <sup>b, c</sup>	0.47 (0.29, 0.65) <sup>d</sup>
<b>Countermovement jump (cm)</b>			
CONTROL	25.9 $\pm$ 4.1	0.5 (-3.5, 4.8)	0.03 (-0.22, 0.29)
PLACEBO	28.7 $\pm$ 5.1	4.4 (2.7, 6.1) <sup>a</sup>	0.23 (0.14, 0.32)
CREATINE	27.3 $\pm$ 5.2	6.5 (3.9, 9.2) <sup>b, c</sup>	0.30 (0.18, 0.41)
<b>20 cm reactive strength index (mm.ms<sup>-1</sup>)</b>			
CONTROL	1.40 $\pm$ 0.6	1.1 (-5.7, 8.3)	0.03 (-0.15, 0.21)
PLACEBO	1.36 $\pm$ 0.4	8.0 (5.6, 10.4) <sup>a</sup>	0.25 (0.18, 0.33)
CREATINE	1.33 $\pm$ 0.3	10.7 (7.5, 14.0) <sup>b, c</sup>	0.42 (0.3, 0.54)
<b>40 cm reactive strength index (mm.ms<sup>-1</sup>)</b>			
CONTROL	1.20 $\pm$ 0.4	4.1 (-5.5, 14.7)	0.14 (-0.2, 0.49)
PLACEBO	1.30 $\pm$ 0.3	10.1 (6.5, 14.0) <sup>a</sup>	0.39 (0.25, 0.53)
CREATINE	1.20 $\pm$ 0.3	13.6 (7.8, 19.7) <sup>b</sup>	0.48 (0.29, 0.68) <sup>d</sup>
<b>Right leg unilateral 20 cm reactive strength index (mm.ms<sup>-1</sup>)</b>			
CONTROL	0.63 $\pm$ 0.3	4.9 (-8.1, 19.7)	0.08 (-0.14, 0.3)
PLACEBO	0.52 $\pm$ 0.2	19.1 (11.5, 27.2) <sup>b</sup>	0.44 (0.28, 0.61)
CREATINE	0.65 $\pm$ 0.3	16.2 (10.0, 22.7) <sup>b</sup>	0.40 (0.25, 0.54)
<b>Left leg unilateral 20 cm reactive strength index (mm.ms<sup>-1</sup>)</b>			
CONTROL	0.43 $\pm$ 0.1	1.8 (-8.8, 13.6)	0.06 (-0.31, 0.43)
PLACEBO	0.42 $\pm$ 0.1	16.1 (11.8, 20.5) <sup>a</sup>	0.47 (0.35, 0.59)
CREATINE	0.43 $\pm$ 0.2	17.9 (12.7, 23.4) <sup>b, c</sup>	0.46 (0.34, 0.59)

<sup>a, b</sup>: denote significant difference pre to post training ( $p < 0.05$  and  $p < 0.01$ , respectively). <sup>c</sup>: denote significant difference with the CONTROL post training ( $p < 0.05$ ); <sup>d</sup>: denote significant greater effect compared to PLACEBO and CONTROL groups.

442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467

**Table 3.** Training effects (with 90% confidence limits) for the running anaerobic sprint test (RAST), change of direction speed, 20-m sprint and endurance performance for the control group (CONTROL, n = 10), plyometric training group receiving placebo (PLACEBO, n = 10) and plyometric training group receiving creatine supplementation (CREATINE, n = 10).

	Baseline Mean $\pm$ SD	Performance change (%)	Effect sizes
<b>RAST mean sprint time (s)</b>			
CONTROL	7.35 $\pm$ 0.5	-0.6 (-3.2, 2.0)	-0.1 (-0.51, 0.31)
PLACEBO	7.08 $\pm$ 0.6	-4.2 (-6.1, -2.3) <sup>a</sup>	-0.48 (-0.70, -0.26)
CREATINE	7.48 $\pm$ 1.0	-5.3 (-7.6, -3.0) <sup>b,c</sup>	-0.55 (-0.80, -0.31) <sup>d</sup>
<b>20-m sprint (s)</b>			
CONTROL	3.99 $\pm$ 0.2	-0.2 (-2.3, 2.0)	-0.05 (-0.58, 0.49)
PLACEBO	3.87 $\pm$ 0.3	-3.2 (-4.4, -2.1) <sup>a,c</sup>	-0.41 (-0.56, -0.27)
CREATINE	3.98 $\pm$ 0.4	-3.3 (-4.6, -2.0) <sup>b,c</sup>	-0.35 (-0.48, -0.21)
<b>Change of direction speed test time (s)</b>			
CONTROL	19.4 $\pm$ 0.8	-0.5 (-2.2, 1.2)	-0.14 (-0.64, 0.35)
PLACEBO	18.8 $\pm$ 1.2	-2.8 (-4.1, -1.6) <sup>a</sup>	-0.46 (-0.67, -0.26)
CREATINE	19.3 $\pm$ 1.1	-2.9 (-3.9, -1.8) <sup>b</sup>	-0.55 (-0.75, -0.34)
<b>20-m multi stage shuttle run test (min)</b>			
CONTROL	7.4 $\pm$ 1.9	2.0 (-1.3, 5.4)	0.06 (-0.04, 0.17)
PLACEBO	7.8 $\pm$ 1.5	6.4 (3.2, 9.6) <sup>a</sup>	0.32 (0.16, 0.47)
CREATINE	8.0 $\pm$ 1.6	7.2 (2.6, 12.1) <sup>b</sup>	0.34 (0.12, 0.56)

<sup>a, b</sup>: denote significant difference pre to post training ( $p < 0.05$  and  $p < 0.01$ , respectively). <sup>c</sup>: denote significant difference with the CONTROL post training ( $p < 0.05$ ). <sup>d</sup>: denote significant greater effect compared to PLACEBO and CONTROL groups.