

1 **An investigation into the daily level of voluntary activity of stabled riding school horses.**

2 H. Brown^{a*}, D. Marlin^b and P. Harris^c

3

4 ^a Hartpury College, Hartpury House, Gloucester, GL19 3BE, UK, Email:
5 hieke.brown@hartpury.ac.uk

6 ^b Science Supplements. Email: dm@sciencesupplements.co.uk

7 ^c Equine Studies Group, WALTHAM Centre for Pet Nutrition, Leicestershire, UK. Email:
8 pat.harris@effem.com

9 * Corresponding author

10

11

12 **Abstract**

13

14 The importance of correct feeding practice has been highlighted by the increase of prevalence
15 of obesity in horses. Human research has suggested that voluntary activity (VA) levels may
16 have the ability to influence digestive energy (DE) requirements, accounting for 15 – 50% of
17 human daily energy expenditure. Therefore, the aim of this study was to investigate whether
18 levels of non-structured exercise differ between stabled horses with similar bodyweight (BW)
19 and similar structured workloads, but with different estimated DE intakes to maintain their
20 BW. Twelve mature horses were selected based on their estimated DE intake and BW, and
21 were paired according to their BW, breed, estimated DE intake, and structured exercise. Within
22 each pair, one horse (L) had a relatively lower estimated DE intake than the other horse (H) to
23 maintain a similar, constant BW. Estimated DE intake was significantly ($P < 0.01$) different
24 between Group L and Group H. Each pair was observed for 72 hours during which structured
25 exercise and non-structured exercise were measured. HR was used as a measure of workload
26 during the structured exercise. Two RT3 accelerometers, located on a roller (RT3-R) and head
27 collar (RT3-H), were used to measure VA levels when stabled in addition to visual observations
28 using focal sampling between 07:00h and 18:00h. RT3-R and RT3-H activity levels were not
29 significantly ($P > 0.05$) different between individual horses. Median activity counts were
30 significantly ($P \leq 0.001$) higher during the day time (06:30h – 18:29h) compared to the night
31 time (18:30h – 06:29h). Both RT3-R and RT3-H were significantly ($P \leq 0.001$) higher around
32 feeding time. Activity measured using RT3 accelerometers suggested higher activity levels for
33 horses in Group H over the 24 hour periods, although this was not significant ($P > 0.05$). It was
34 therefore concluded, that differences in VA levels during stabling could not explain the
35 difference in estimated DE requirements between horses with a similar BW and workload.

36

37 **Keywords**

38 Activity, equine, estimated energy requirement, accelerometer

39

40

41

42 1.0 Introduction

43 The prevalence of obesity in horses used for leisure has been increasing, potentially as a direct
44 result of domestication and current management practices (Argo, 2009; Sillence *et al.*, 2002;
45 Wyse *et al.*, 2008). Weight gain is likely to occur when there is an imbalance between energy
46 intake via the diet and energy requirement (Geor & Harris, 2005; Harris and Kronfeld, 2003).
47 Various intrinsic and extrinsic factors have the potential to affect the digestible energy (DE)
48 requirements of an individual horse being exercised including the horse's body weight (BW)
49 and age plus any individual absorptive and metabolic factors, stage of training and workload,
50 ability and weight of the rider, as well as environment and terrain (Anon, 2007; Harris, 1997;
51 Hintz and Cymbaluk, 1994). However, it has been suggested that the small individual
52 differences in workload between individuals of a similar bodyweight, age and fitness
53 undertaking similar types of structured exercise under the same environmental conditions
54 cannot always explain the variability in estimated DE intake required to maintain a constant
55 BW (Dekker *et al.*, 2007; Dekker, 2009). Individual differences in absorptive capacity or
56 metabolic efficiency are a possible explanation (Anon, 2007), but other less obvious
57 differences may exist between individuals including the amount of non-structured exercise they
58 undertake.

59 Within current management systems, horses are often housed in individual stables with limited
60 or no turn-out into a paddock or field, which limits their ability to undertake free movement or
61 non-structured exercise (Brehme and Rose, 2007; Harris, 1999; Petersen *et al.*, 2005). In fact
62 very few studies have focussed on the levels of non-structured activity or voluntary activity
63 (VA: any movement within the stable or during turn-out that is not a result of human
64 encouragement) of horses during stabling or turn-out (Rose-Meierhofer *et al.*, 2010; Werhahn
65 *et al.*, 2012) and the potential influence of this on individual DE requirements. Human studies
66 have suggested that VA may play an important role in the regulation of energy balance, where
67 VA is associated with posture maintenance, fidgeting and all other daily activities which are
68 not classified as exercise (Levine *et al.*, 1999; Levine *et al.*, 2000). Reported data in humans
69 has suggested that VA can account for ~15% of the daily exercise expenditure in sedentary
70 individuals to more than 50% of the daily energy expenditure in extremely active individuals
71 (Deriaz *et al.*, 1992; Levine, 2003; Livingstone *et al.*, 1991; Ravussin *et al.*, 1986), and may
72 contribute to the inter-individual variation in susceptibility to weight gain as a response to
73 overfeeding (Levine *et al.*, 1999).

74 The aim of this study was therefore to investigate whether VA levels differed between stabled
75 horses with similar BW, undertaking similar levels of structured exercise but requiring different
76 estimated DE intakes to maintain their BW. It was hypothesised that VA levels would differ
77 during a 24 hour period and that horses with a higher estimated DE intake would have higher
78 levels of VA.

79

80 **2.0 Material and Methods**

81 *Horses and diets*

82 Twelve mature horses (8 geldings and 4 mares) with mean age of 10.9 ± 4.0 years (range: 5 –
83 18 years) of mixed breeds were selected from a large population of experienced riding school
84 horses (Table 1). All horses were on a routine anthelmintic and dental care program and only
85 animals in good health and dental status, which did not exhibit any stereotypic behaviour, were
86 recruited. All horses had similar levels of structured exercise. This exercise consisted of on
87 average ten student training sessions per week (range: 8 – 12 sessions per week), which
88 included flat work and low level jumping. During the weekend, horses were hacked out or
89 walked out. None of the horses were turned out during the study period. All horses received
90 their standard commercially available diet during the study. The compound feeds were
91 provided twice daily at 07:00h and 16:30h and the hay or haylage was provided three times
92 per day at 07:30h, 12:30h, and 17:00h.

93

94 **Table 1. Mean (\pm SD) age, height, bodyweight (BW), body condition score (BCS),**
95 **estimated DE intake and relative workload (RW) for horses in group L (n = 6) and horses**
96 **in group H (n = 6).** Different letters (a, b, c etc.) indicate significant differences ($P < 0.05$)
97 between the two groups, i.e. ‘a’ is significantly different from ‘b’, etc.

98

Variable	Group L (n = 6)	Group H (n = 6)
Age (years)	10.3 ± 4.9 ^a	11.5 ± 3.3 ^a
Height (cm)	157.7 ± 1.0 ^a	162.0 ± 4.6 ^a
BW (kg)	518.8 ± 25.8 ^a	530.0 ± 30.7 ^a
BCS	4.32 ± 0.79 ^a	4.32 ± 0.81 ^a
DE intake (MJ/day)	102.3 ± 13.9 ^a	125.3 ± 8.8 ^b
RW (%HR _{max})	42.0 ± 4.2 ^a	41.6 ± 3.8 ^a

100

101 Horses were selected based on their estimated DE intake and their BW, which was monitored
102 weekly (Eziweigh 2, Tru-Test Ltd, Auckland, New Zealand) for a five week period prior to the
103 start to confirm that the chosen horses were maintaining their BW on their individual diets. A
104 BW variation of 1% BW between weekly weighing sessions was considered insignificant and
105 the result of daily variation due to last time of defecation or urination. In addition, body
106 condition (BC) of the horses was determined using either Henneke *et al.* (1983) or Kienzle and
107 Schramme (2004) dependent on the horse's breed. The individual DE intakes were estimated
108 based on the DE values (MJ/kg) of commercially available feeds as provided by the feed
109 manufacturers, and the DE values of hay or haylage as determined by an independent
110 laboratory (Direct Laboratories, Wolverhampton, UK).

111 Horses were paired based on a number of variables, including BW, breed, structured exercise
112 load, and estimated DE intake. A maximum difference of 6% BW between paired horses was
113 allowed. Each pair consisted of one horse (L) which received a relatively lower amount of
114 estimated DE (MJ/kg) (mean 18 ± 8% lower) than the other horse (H) in order to maintain a
115 similar constant BW.

116

117

118 *Data collection*

119 Two RT3 accelerometers (StayHealthy Inc, California, USA), located on an anti-cast roller
120 (RT3-R) and on an easy-snap head collar (RT3-H), were used to determine VA levels. Previous
121 work (Dekker, 2009) had demonstrated that these positions provided the most reliable
122 measurements. Both accelerometers were set to record activity over one-minute intervals.

123 Each pair of horses was observed over a period of 72 hours. During this period, horses were
124 used for their normal riding sessions during which heart rate (HR) was monitored (Polar S610i,
125 Polar, Kempele, Finland), but RT3 accelerometers were removed. Relative workload (RW)
126 was calculated as the mean HR during work as a percentage of the horse's estimated maximum
127 HR (HR_{max}). HR_{max} was estimated using the equation: $HR_{max} = 224 - \text{age}$ (Vincent *et al.*,
128 2006).

129 In addition to data collected by the RT3 accelerometers, horses were observed visually between
130 07:00h and 18:00h during the 72h study period. Focal sampling with continuous recording was
131 used for 30 minutes, with one horse being the focus at any one time. Observation periods were
132 allocated evenly between paired horses, enabling a representation of their activity pattern to be
133 evaluated. Horses were observed before, during, and after concentrate meals as well as between
134 meals and training sessions. The observer was situated outside the stable with a clear view of
135 the horse. Different stable activities and behaviours were recorded during one minute time
136 periods, which coincided with the RT3 accelerometers' intervals. These activities were further
137 categorised as leg movement, head movement, and total movement per minute during data
138 analysis.

139

140 *Statistical analysis*

141 Data were analysed using SPSS version 15.0 for Windows. Using the Shapiro-Wilk W test it
142 was determined that raw data was not normally distributed ($P > 0.05$). Therefore, the median of
143 the data was used when aggregating the rough data to one hour intervals ($n = 288$), time periods
144 (day: 06:30h – 18:29h and night: 18:30h – 06:29h, $n = 24$), day level ($n = 36$) and horse level
145 ($n = 12$). Spearman's correlation coefficient (r_s) with a significance level of $P \leq 0.05$ was used
146 to determine correlations between monitored activity and observed activity using the raw data.

147 Aggregated data for hourly intervals, day periods, day level, and horse level showed a normal
148 distribution established using the Shapiro-Wilk W test with a significance level of $P \leq 0.05$,
149 allowing for the use of further parametric tests. Pearson's correlation coefficient (r) was used
150 to determine correlations between horse characteristics and aggregated data at horse level.
151 Secondly, a one-way ANOVA with a Bonferroni post-hoc test with a significance level of
152 $P \leq 0.05$ was used to determine whether VA levels were consistent during 72h observation
153 period using aggregated data at day level, as well as whether the mean daily activity measured
154 by RT3-R and RT3-H were significantly different between horses. Thirdly, an independent t-
155 test with significance level of $P \leq 0.05$ was used to determine significant differences for RT3-R
156 and RT3-H between day time and night time using the aggregated data for time periods.
157 Finally, aggregated data at hour level was used to determine patterns of daily activity in
158 addition to observed activity through visual observations.

159

160 **3.0 Results**

161 Throughout the study the horses maintained a consistent BW (mean variation 2.2 ± 2.3 kg BW)
162 and BCS and remained clinically healthy. All horses remained sound and maintained their
163 normal weekly workload. Within pairs the average difference in BW was 3.3 ± 1.7 %BW.

164

165 *Observed activity*

166 On average each individual horse was observed for 515.6 ± 94.4 minutes (range: 451 – 627
167 minutes) between 07:00h and 18:00h over the three consecutive days. The number of observed
168 minutes was not significantly different ($P > 0.05$) between horses in Group L and Group H
169 (506.8 ± 93.5 minutes and 524.3 ± 103.3 minutes, respectively). The main observed activities
170 included standing ($52.4 \pm 10.1\%$), eating hay ($41.7 \pm 9.6\%$) and eating concentrate feed (5.8%
171 $\pm 2.6\%$). Observed activity showed weak, but significant ($P < 0.05$) correlation with monitored
172 activity. Activity measured by RT3-R showed a weak correlation ($r = 0.22$) with the amount
173 of time spent walking and moving around the stable. The activity recorded by the RT3-H
174 showed a moderate correlation with eating ($r = 0.40$) and vertical head movement ($r = 0.24$).


175

176

177 *Monitored daily activity*

178 Data from RT3-R (median counts = 6.5, range: 0 – 144) was significantly ($P < 0.0001$) lower
179 compared to data from RT3-H (median counts = 144, range: 0 – 2073.5). There was no
180 significant difference between individual horses for median daily activity as measured by the
181 RT3-R ($P > 0.05$) and RT3-H ($P > 0.05$). No significant correlations ($P > 0.05$) were found
182 between median activity counts of the individual horses and the horses' characteristics,
183 including age, BW, BCS, DE intake, and RW.

184




185

186 **Figure 1. Median RT3 roller and median RT3 head activity during the day time (06:30h**
187 **– 18:29h) and the night time (18:30h – 06:29h) (N = 12). N equals the number of horses.**
188 Bars with different letters are significantly different ($P < 0.05$), i.e. 'a' is significantly different
189 from 'b', etc.

190

191 Median activity counts of the horses was significantly ($P \leq 0.001$) higher during the day time
192 (06:30h – 18:29h) for both RT3-R and RT3-H compared to night time (18:30h – 06:29h)
193 (Figure 1). In addition, RT3-R and RT3-H differed significantly between individual hours
194 (Figure 2), with increased activity being measured around 07:00h, 12:00h, and 16:30h. This
195 was associated with the time of feeding. Both RT3-R and RT3-H were significantly ($P \leq 0.001$)
196 higher during the eating of compound feeds compared to the eating of hay.

197



198

199 **Figure 2. Activity pattern of mean RT3 roller activity and mean RT3 head activity over**
200 **a 24h period (N = 12).** N equals the number of horses. Arrows indicate occurrence of feeding

201

202 *Activity within pairs*

203 Group L and Group H did not differ significantly ($P > 0.05$) in mean age, mean height, mean
204 BW, mean BCS, and mean RW. Although the use of compound feeds did not differ between

205 the two groups, the estimated DE intake (MJ/day) was significantly ($P < 0.01$) different (see
 206 table 1) as expected. Horses in group L were visually observed to move significantly ($P < 0.05$)
 207 more per minute than horses in group H (12.9 ± 7.8 movements per minute and 11.3 ± 7.2
 208 movements per minute, respectively). However, activity measured using RT3-R and RT3-H
 209 indicated higher overall activity counts for horses in group H compared to group L during the
 210 daytime and night time periods as well as over 24 hour periods, although these differences
 211 were not significant ($P > 0.05$) (Table 2).

212

213 **Table 2. Mean (\pm SD) counts of the RT3 roller and RT3 head activity for horses in Group**
 214 **L (n = 6) and Group H (n = 6).** Mean values for daytime (06:30h – 18:29h), night time (18:30
 215 – 06:29h), and 24 hour period derived from data collected over a period of 72 hours. Different
 216 letter (a, b, c, etc) indicate significant differences ($P < 0.05$) between the two groups, i.e. ‘a’ is
 217 significantly different from ‘b’, etc.

218

Time	RT3	Group L (n = 6)	Group H (n = 6)	All horses (n = 12)
Day time (06:30h – 18:29h)	Roller	20.0 ± 12.8^a	20.5 ± 16.6^a	20.3 ± 14.2
	Head	376.0 ± 147.8^a	322.4 ± 109.4^a	349.2 ± 127.1
Night time (18:30h – 06:29h)	Roller	2.1 ± 3.1^a	5.8 ± 7.0^a	3.9 ± 5.5
	Head	36.8 ± 30.7^a	69.8 ± 47.7^a	53.3 ± 42.0
24 hour period	Roller	9.7 ± 6.3^a	11.3 ± 9.7^a	12.1 ± 13.4
	Head	117.4 ± 113.5^a	184.0 ± 88.9^a	201.3 ± 177.2

219

220 4.0 Discussion

221 The use of accelerometers to objectively measure activity in horses has been reported in various
 222 different contexts (Berger *et al.*, 1999; Burla *et al.*, 2014; Piccione *et al.*, 2008; Scheibe *et al.*,
 223 1998), although to the authors’ knowledge no link has been made to energy expenditure and

224 DE requirements of the horse. With the knowledge that VA levels can account for more than
225 50% of the daily energy expenditure in extremely active human individuals (Deriaz *et al.*, 1992;
226 Levine, 2003; Livingstone *et al.*, 1991; Ravussin *et al.*, 1986), it could be presumed that similar
227 principles could be underlying to the anecdotal evidence that horses of similar BW and
228 structured workload have significantly different DE requirements

229 Because of the various variables known to influence DE requirements in horses, the horses in
230 this study had been paired according to a similar BW, age, breed, and estimated RW, but with
231 significantly ($P<0.05$) different DE requirements (MJ/day) to maintain their BW and RW.
232 Breed was considered an important factor, as previous results have indicated that reactivity,
233 temperament, and VA levels tend to vary between different breeds (Hausberger and Muller,
234 2002; Lloyd *et al.*, 2008; Søndergaard and Ladewig, 2004). Although horses in Group L were
235 visually observed to move significantly ($P<0.05$) more than horses in Group H, this was not
236 mirrored in the recordings of the RT3-R and the RT3-H activity. Therefore, it can be presumed
237 that other variables have a larger influence on the daily DE requirements of horses than VA
238 levels in the stable.

239 Although VA levels were similar over the 72h observation periods, day time activity (06:30h
240 – 18:29h) was found to be significantly higher ($P\leq 0.001$) compared to night time activity
241 (18:30h – 06:29h) for all horses. The higher activity levels recorded during the day time could
242 be attributed especially to behaviour around feeding time (Figure 2). Increased activity levels
243 have previously been recorded as anticipation behaviour (Cooper and Mason, 1998; Cooper *et al.*
244 *et al.*, 2000; Cooper *et al.*, 2005). However, in contrast to these studies, it is important to note that
245 none of the horses in the current study performed any stereotypical behaviours. It can therefore
246 be suggested that anticipation of feeding can result in an increase in activity in horses in
247 general, not just those with stereotypies. Day time activity did not differ significantly between
248 the two groups (Table 2), suggesting further that activity on the yard and the horses' exercise
249 and feeding regimen strongly influenced day time VA levels. Indeed, horses have previously
250 been shown to spend significantly less time resting after exercise and more time feeding and
251 drinking (Caanitz *et al.*, 1991).

252 Although no visual observations were undertaken during the night time, it may be presumed
253 that horses spent the majority of their time resting and ingesting hay, as reported by Greening
254 *et al.* (2013). It is of interest to note that VA levels were slightly higher for horses in Group H,
255 although the difference was not significant ($P>0.05$). Horses were all stabled on a combination

256 of rubber matting and shavings, limiting the variable of bedding on behaviour and activity
257 patterns during the night time period (Greening *et al.*, 2013). The difference between the two
258 groups may potentially be the result of large individual differences within a relatively small
259 sample size. Therefore, it is suggested that a larger sample size would be required to ascertain
260 this. Although VA levels during turn-out were not evaluated during this study, this should also
261 be considered as an influencing factor for differences in daily DE requirements between similar
262 horses. It is suggested that further research is undertaken within this area.

263

264 **5.0 Conclusion**

265 The present study objectively measured the activity levels of stabled, general purpose horses
266 with no stereotypical behaviours. Differences in VA levels could not explain the difference in
267 estimated DE requirements between pairs of animals of a similar age with similar BW,
268 workload, etc., although the trend for differences over night might suggest that it potentially
269 could have a small role to play in stabled horses.

270

271 **Acknowledgements**

272 We would like to acknowledge Hartpury College and SPILLERS for the funding of this project
273 and would like to thank the yard staff, riding instructors and students at Hartpury College for
274 their help during this study.

275

276 **References**

277 Anon, 2007. Nutrient requirements of horses, 6th edition. National. Academic Press,
278 Washington DC, USA.

279

280 Argo, C. McG., 2009. Appraising the portly pony: Body condition scores and adiposity. *The*
281 *Veterinary Journal* 179: 158-160

282

283 Berger, A., Scheibe, K.M., Eichhorn, K., Scheibe, A. and Streich, J., 1999. Diurnal and
284 ultradian rhythms of behaviour in a mare group of Przewalski horse (*Equus ferus przewalskii*),
285 measured through one year under semi-reserve conditions. Applied Animal Behaviour Science
286 64: 1-17

287

288 Brehme, U. and Rose, S., 2007. Effect of different activity and space offers on the activity
289 behaviour of stallions. Landtechnik. 62: 408-409

290

291 Burla, J-B., Ostertag, A., Schulze Westerath, H. and Hillmann, E., 2014. Gait determination
292 and activity measurement in horses using an accelerometer. Computers and Electronics in
293 Agriculture 102: 127-133

294

295 Caanitz, H., O'Leary, L., Houpt, K., Petersson, K. and Hintz, H., 1991. Effect of exercise on
296 equine behavior. Applied Animal Behaviour Science 31: 1-12

297

298 Cooper, J.J. and Mason, G.J., 1998. The identification of abnormal behaviour and behavioural
299 problems in stabled horses and their relationship to horse welfare: a comparative review.
300 Equine Veterinary Journal Supplement 27: 5-9

301

302 Cooper, J.J., McDonald, L. and Mills, D.S., 2000. The effect of increasing visual horizons on
303 stereotypic weaving: implications for the social housing of stabled horses. Applied Animal
304 Behaviour Science 69: 67-83

305

306 Cooper, J.J., McCall, N., Johnson, S. and Davidson, S., 2005. The short-term effects of
307 increasing meal frequency on stereotypic behaviour of stabled horses. Applied Animal
308 Behaviour Science 90: 351-364

309

310 Dekker, H., 2009. A study of the effects of diet and management on the activity and body
311 condition of stabled horses. PhD thesis, University of the West of England

312

313 Dekker, H., Marlin, D., Alexander, L., Bishop, R., and Harris, P., 2007. A pilot study
314 investigating the relationship between perceived and actual workload and estimated energy
315 intake in riding centre horses. *Equine and Comparative Exercise Physiology* 4: 7-14

316

317 Deriaz, O., Fournier, G., Tremblay, A., Despres, J.P. and Bouchard, C., 1992. Lean-body-mass
318 composition and resting energy expenditure before and after long-term overfeeding. *The*
319 *American Journal of Clinical Nutrition* 56: 840-847

320

321 Geor, R.J. and Harris, P.A., 2005. Nutritional Management of Endurance Horses. In: Harris,
322 P.A., Mair, T.S., Slater, J.D. and Green, R.E. (Eds.), *Equine Nutrition for All. The 1st BEVA*
323 *and WALTHAM Nutrition Symposia*. *Equine Veterinary Journal*, Newmarket, UK, pp. 71 - 78

324

325 Greening, L., Shenton, V., Wilcockson, K. and Swanson, J., 2013. Investigating duration of
326 nocturnal ingestive and sleep behaviors of horses bedded on straw versus shavings. *Journal of*
327 *Veterinary Behaviour* 8: 82-86

328

329 Harris, P.A., 1997. Energy sources and requirements of the exercising horse. *Annual Review*
330 *of Nutrition* 17: 185-210

331

332 Harris, P.A., 1999. Review of equine feeding and stable management practices in the UK
333 concentrating on the last decade of the 20th century. *Equine Veterinary Journal Supplement* 28:
334 46-54

335

336 Harris, P.A. and Kronfeld, D.S., 2003. Influence of dietary energy sources on health and
337 performance. In: Robinson, N.E. (Ed.), Current Therapy in Equine Medicine 5. WB Saunders,
338 Philadelphia, USA, pp.698-704

339

340 Hausberger, M. and Muller, C., 2002. A brief note on some possible factors involved in the
341 reactions of horses to humans. Applied Animal Behaviour Science 76: 330-344

342

343 Henneke, D.R., Potter, G.D., Kreider, J.L. and Yeates, B.F., 1983. Relationship between
344 condition score, physical measurement and body fat percentage in mares. Equine Veterinary
345 Journal 15: 371-372

346

347 Hintz, H.F. and Cymbaluk, N.F., 1994. Nutrition of the horse. Annual Review of Nutrition 14:
348 243-267

349

350 Kienzle, E. and Schramme S.C., 2004. Beurteilung des Ernährungszustandes mittels Body
351 Condition Scores und Gewichtsschätzung beim adulten Warmblutpferd. Pferdeheilkunde 20:
352 517-524

353

354 Levine, J.A., Eberhardt, N.L. and Jensen, M.D., 1999. Role of nonexercise activity
355 thermogenesis in resistance to fat gain in humans. Science 283: 212-214

356

357 Levine, J.A., Schleusner, S.J. and Jensen, M.D., 2000. Energy expenditure of nonexercise
358 activity. The American Journal of Clinical Nutrition 72:1451-1454

359

360 Levine, J.A., 2003. Non-exercise activity thermogenesis. Proceedings of the Nutrition Society
361 62: 667-679

362

363 Livingstone, M.B.E., Strain, J.J., Prentice, A.M., Coward, W.A., Nevin, G.B., Barker, M.E.,
364 Hickery, R.J., McKenna, P.G. and Whitehead, R.G., 1991. Potential contribution of leisure
365 activity to the energy expenditure patterns of sedentary populations. *British Journal of Nutrition*
366 65: 145-155

367

368 Lloyd, A.S., Martin, J.E., Bornett-Gauci, H.L.I. and Wilkinson, R.G., 2008. Horse personality:
369 variation between breeds. *Applied Animal Behaviour Science* 112: 369-383

370

371 Petersen, S., Tolle, K.H., Blobel, K.J., Grabner, A. and Krieter, J., 2005. Evaluation of horse
372 keeping in Schleswig-Holstein. *In: 56th Annual Meeting of the European Association of*
373 *Animal Production, Uppsala, Sweden* 33. EAAP – Book of Abstracts No 11

374

375 Piccione, G., Costa, A., Giannetto, C, and Caola, G., 2008. Daily rhythms of activity in horses
376 housed in different stabling conditions. *Biological Rhythm Research* 39: 79-84

377

378 Ravussin, E., Lillioja, S., Anderson, T.E., Christin, L. and Bogardus, C., 1986. Determinants
379 of 24-hour energy expenditure in man. Methods and results using a respiratory chamber. *The*
380 *Journal of Clinical Investigation* 78: 1568-1578

381

382 Rose-Meierhofer, S., Klaer, S., Ammon, C., Brunsch, R. and Hoffmann, G., 2010. Activity
383 behaviour of horses housed in different open barn systems. *Journal of Equine Veterinary*
384 *Science* 30: 624-634

385

386 Scheibe, K.M., Schleusner, Th., Berger, A., Eichhorn, K., Langbein, J., Dal Zotto, L. and
387 Streich, W.J., 1998. ETHOSYS ® -new system for recording and analysis of behaviour of free-
388 ranging domestic animals and wildlife. *Applied Animal Behaviour Science* 55: 195-211

389

390 Sillence, M., Noble, G. and McGowan, C., 2002. Fast food and fat fillies: the ills of western
391 civilization. *The Veterinary Journal* 172: 396-397

392

393 Søndergaard, E. and Ladewig, J., 2004. Group housing exerts a positive effect on the behaviour
394 of young horses during training. *Applied Animal Behaviour Science* 87: 105-118

395

396 Vincent, T.L., Newton, J.R., Deaton, C.M., Franklin, S.H., Biddick, T., McKeever, K.H.,
397 McDonough, P., Young, L.E., Hodgson, D.R. and Marlin, D.J., 2006. Retrospective study of
398 predictive variables for maximal heart rate (HR_{max}) in horses undergoing strenuous treadmill
399 exercise. *Equine Veterinary Journal Supplement* 36: 146-152

400

401 Werhahn, H., Hessel, E.F., Herman and Van de Weghe, H.F.A., 2012. Competition horses
402 housed in single stalls (I): Behavior and activity patterns during free exercise according to its
403 configuration. *Journal of Equine Veterinary Science* 32: 45-52

404

405 Wyse, C.A., McNie, K.A., Tannahil, V.J., Murray, J.K. and Love, S., 2008. Prevalence of
406 obesity in riding horses in Scotland. *The Veterinary Record* 162: 590-591

407

408